
EXAMINATION OF THE EFFECT OF SOLID MINERAL MINING (LEAD) ON THE WATER RESOURCES IN ABAKALIKI, EBONYI STATE

CHUKWU, KEVIN EJIKE Ph.D.

Department of Geography and Meteorology,
Faculty of Environmental Sciences
Enugu State University of Science and Technology (ESUT), Enugu State

Abstract

Lead mining activities in Ishiagu have drastically affected the groundwater resource of the area, contaminating it with significant concentrations of metallic lead and zinc. Season affects lead concentrations adversely with dry season recording higher concentration values. Heavy metals (lead and zinc) distribution in the groundwater and fishes, follow the same pattern, indicating a close correlation between the levels in groundwater and in fishes. The fact that both dry and rainy season mean levels of lead exceeded WHO (1985) and NAFDAC (2006) recommended limits in food and water, suggested that both groundwater and fishes of the area are endemic for human consumption. Based on that, metal pollution in Ishiagu should be monitored closely due to possible health implication to consumers of the fishes and groundwater users in the area.

Keywords: Solid Mineral, Mining, Lead, Water Resources, Abakaliki

Introduction

Lead-Zinc ores are usually found together. They are often associated with copper and silver. Lead-zinc ores are used in the production of batteries, electrical cables, solders glass and even protective coatings for other metals. Mining industries are now viewed as key drivers of economic growth and the development process (Bradshaw, 2005), and as lead sectors that drive economic expansion which can lead to higher levels of social and economic well-being (Bridge, 2008).

Mineral is an inorganic substance found in nature with a characteristics chemical composition and in some cases, of definite crystal structure. In the absence of adequate portable water supply majority of residents depends on surface water streams namely Onuafia, Ihekoyi and Akpaudo, abandoned Zinc and lead Ores mining pits, and Ugwuadu spring for agricultural and domestic uses respectively. Geologically, the Ishaiagu Lead deposits form part of the Abakaliki Lead field (BESL. 2004). This field occurs within a substantial part of the largest cretaceous sedimentary

basis in Nigeria known as the Benue trough. The Ishiagu mineral belt is part of the tectonic – mineralization zone forming the lower Benue field.

Heavy metals are chemical elements with specific gravity (density) of at least five times that of water (Liprot, 1983). Heavy metals come from various sources; naturally occurring mineral deposits like lead, zinc, iron ore, marble limestone and chromium (Obienusi, 2004). Industrial processes such as mining activities are other sources of heavy metals. It is true that some metals are very useful to man and manufacturing industries, they are at the same time very toxic to man even at a low concentrations when they enter the aquatic environment.

The pollution of the aquatic environment with heavy metals has become a worldwide problem during recent years, because they are indestructible and most of them have toxic effects on organisms (MacFarlane and Burchett, (2000).

Among the environmental pollutants, metals are of particular concern, due to their potential toxic effect and ability to bioaccumulation in aquatic ecosystems, Censi, *et al*, (2006). The accumulation of heavy metals in freshwater ecosystem has been a major concern. Heavy metals generally enter the aquatic environment through natural (atmospheric deposition, erosion of geological matrix) or anthropogenic activities (caused by industrial effluent, domestic sewage, mining and agriculture wastes) Connell, and Miller (1984) and Vautukuru (2005).

Study Area

Ishiagu is located in the southern part of Ebonyi State where open cast mining of Lead (Pb) and Zinc (Zn) is common. It lies between latitude 7030` and 7037` N and longitude 5052` and 6000` E. Ishaigu is richly endowed with mineral deposits, which are mainly heavy metal sulfide ores, mostly galena(PbS). It is intensively and competitively exploited by several mining industries for over thirty (30) years now. Ishiagu is made up of eleven (11) villages with five active mining villages, namely: Amagu, Amaonye, Ihetutu and Ameze. It is surrounded by Afikpo to the North, LokpaNta to the East, Lekwesi to the West and Uburu to the South (Figure 1) and the study locations shown in Figure 2.






Fig. 1: Map of Ebonyi State showing Ivo L.G.A (Survey Dept Abakaliki, 2012).




Fig. 2: Map of Ishiagu Showing Study Locations (Survey Department, Abakaliki, 2012).

Importance of Lead Mining

It is estimated that Nigeria has over 10,000,000 tons of lead ore deposit and it is an important component in car battery production. The prospect of global economy is promising, so the automotive industry which is closely related to the lead market, is witnessing growth thereby creating more demands for the product. Lead-zinc ores are used in the production of batteries, electrical cables, solders glass and even protective coatings for other metals

Effect of Lead Mining on Water

Socio-economically the people of Ishiagu are mainly farmers. The crops cultivated include yams, cassava and rice. Cassava and yam are commercially produced and therefore form the economic base of the area. In the absence of adequate portable water supply majority of residents depends on surface water streams.



The mineral assemblage and associations in the area comprise both primary and secondary mineralization (Offordile 2001). The primary metalliferous deposits in the area include Galena, Sphalerite, Chaleopyrite, Marcasite, Cerrurite and Pyromorphite while the gangue minerals associated with the metallic ores include siderite, calcite, fluorite, barite and quartz. Successive mining activities by various companies have depleted this reserve over the years.

Pollution of surface water streams and spring used for agricultural and domestic purposes, as well as, rising noise and suspended air particulate levels across the Ishiagu communities. The presence of large excavated pits used for zinc and lead ores exploration/mining (without any form of backfilling as earlier indicated in Environmental Management Plans (EMP) by the various mining operators and its attendant land damage and degradation) has forced farmers to source farmlands at far distances from their communities of domicile.

Furthermore, the pollution of Onuafia, Ihekoyi and Akpaudo streams and, in particular that of Ugwuado spring used for drinking purpose by the agrarian settlements (in an environment where amenities and services such as clean water and health care are grossly inadequate is to say the least undesirable (Essaghah, Ogbona and Ugwuanyi 2006).

With the current levels of Suspended Particulate Matter (SPM) in all locations well above Federal Ministry of Environment and World Health Organization (WHO) standards (during the dry season), there is no doubt that the health status of these agrarian communities have become worse for it. Obviously policy has a role to play in reversing these negative socioeconomic and healthcare indications.

Air Quality

The concentration of gaseous pollutants were very low, particularly those of hydrogen sulphide and carbon monoxide. However, the concentration of Suspended Particulate Matter (SPM) was much higher which could be attributed to dust haze.

Air pollutants are intimately associated with a number of respiratory diseases, largely in aggravating existing disorders such as emphysema, bronchitis and asthma (Ardayfio-schandoof and Asiedu, 2003).

Loss of natural vegetation possess a risk for the surrounding area because the absence of vegetation facilitates lateral wind erosion of metal contaminated particles, and may enhance the



volume of water penetrating through the soil and eventually reaching and contaminating the underlying ground water (Vangronsveld et al 1991; Vangronsveld et al 1995b). These consequences are undesirable in a fragile, rural, agrarian settlements sustained in large part by agriculture, and where amenities and services such as clean water and health care are grossly inadequate.

Conclusion

The environmental impact of solid mining activities in Ishiagu, a community known for large deposits of lead and zinc ores, has potential of becoming an industrial area in Ebonyi State of Nigeria. One way of ensuring continued exploitation of valuable minerals and harness their positive contributions for sustainable economic and physical transformation of the local communities is to protect existing environmental systems and resources that support the agricultural pursuits and livelihood of the people (Essaghah 2010a; Essaghah and Alabi 2013).

Recommendations

In view of the adverse effect of lead mining on water resources in Ishiagu and the health implications of drinking contaminated water, the following measures are recommended to guard against the future and further drinking of such water and its deadly effect.

- ❖ Ebonyi State Government through the Ministries of Agriculture, Environment and Water resources should provide direct support services to the people in the area of rural water supply projects including distribution of subsidized water treatment chemicals/kits, fertilizer distribution and other farm implements without delay. This will help to amend the difficulties faced by the rural farmers and help arrest the rising incidence of dysentery, typhoid and respiratory related disorders and or diseases.
- ❖ Environmental Regulatory Agencies should stipulate specific mitigation measures that industry and mining companies must implement in their host communities before granting approval permits. Rather than leave such decisions to individual companies. Regulatory agencies should embark on the development of data base on the problems resulting from industrial operations in the implementation of mitigation measures by investors in these areas.



- ❖ The Ebonyi State Environmental Protection Agency (EBSEPA) should rise up to its statutory responsibilities (in collaboration with the Ebonyi State Ministry of Environment) by embarking on effective monitoring of the activities of mining and quarrying industries in the state to ensure that industrial operators fulfill their social responsibilities to the communities where they operate.
- ❖ The Federal Government should adopt a deliberate policy of direct intervention in the provision of necessary social amenities in industrial areas to complement the efforts of industrial operators if its current emphasis on industrialization and employment creation is to yield desirable results. This will go along way in complementing the various residual mitigation measures usually adopted by mining companies in their areas of operation (Essaghah and Ugwuayi, 2006).

Until this is done industrial concerns in general and mining operators in particular will continue to smile to banks at the expense of the very poor rural farmers' land; teaming inhabitants who bear the impact of the damages caused by them and at whose instance the government encourages industrialization.

References

- American Public Health Association (APHA) (1995). Standard Methods for the Examination of Water and Waste Water, APHA. Washington D.C.
- Ardayfio-schardoff .E. and Asizdu, A.B.(2003). The Tropical Environment and Environmental Pollution: The Need for Action. In E.N Adinna, O.B. Ekop and V.I Attah (Eds): Environmental Pollution and Management in the Tropics. SNAAP Press Limited Enugu.
- Association of Official Analytical Chemists (AOAC) (1974). Chemical Analytical Methods AOAC Washington D.C
- Bradshaw. M. J. (2005). Population Resources, Development and the Environment, In: Daniels P.E. et al eds. An Introduction to Human Geography: Issues for the 21st Century. 2nd Edition.
- Bastida .E. (2001). A Review of the Concept of Security of Tenure: Issues and Challenges. Available at <http://www.dundee.ac.uk/cepmlp/journal/htm/vol7./article7-17 htm> (2010)



- Bridge. G. (2008) Economic Geography Natural Resources; In: International Encyclopaedia of Human Geography. (Eds). Kitchen and Thrift. Elsevier
- Built Environmental Services Limited (BESL),(2004). Environmental Impact Assessment of Mining of Lead and Zinc Ores in ML 17439 Ishiagu BESL, Enugu.
- Derome, J AND Lindroos J (1998).Effects of Heavy Metal Accumulation on Macronutrient Availability and Acidification Parameters in the Vicinity of the Hargavaltta Cu-Ni Smelter SW. Finland. *Environmental Pollution* 99,225-32.
- Essaghah A.A.E. and Alabi, M.O. (2013). Strategy for Promoting Effective Management of Environmental Effects from Brewery Industries at the 9th Mile Corner Industrial District of Enugu State. *International Journal of Energy and Environmental Research: Vol. 3 No. 1*.
- Essaghah, A.A.E. (2011). A Strategy for Managing Environmental Hazards at the Community Level in Nigeria. Bring a LEAD Paper Presented during the 1st Natural Annual General Meeting (AGM) of the Environmental Management Association of Nigeria (EMAN) held on 7th May 2011 at Sunshine Hotel, Ebeano Estate. New Haven Enugu metropolis, Nigeria.
- Essaghah, A.A.E, Ogbonna .C. and Ugwuanyi B.C. (2006). Lead and Zinc Ores Mining and Environmental Degradation in Ishiagu Community of Ebonyi State. Being a Paper delivered during the Annual General National Conference of Nigerian Environmental Society (NES) held at the Presidential Hotel Enugu between 7th – 9thDecember 2006.
- Essaghah, A.A.E and Ugwu, L.N. (2008). Management Strategy for Monitoring and Auditing Environmental Effects of Brewery Industries in Nigeria: The case of Nigeria Bottling Company Plc. *ESUT Journal of Politics, Ecology and Environment.Vol.1. No.1*
- Essaghah, A.A.E. (2010). The Menace of Flooding in a Rapidly Urbanizing Environments: A case study of Warri Metropolis. Being A paper Delivered on the 4th Edition of Late Professor F.M.A
- Esteves, A. (2008). Mining and Social Development: Refocusing Community Investment and Using Multicriteria Decision Analysis. *Res Policy* 33.39-47
- Girones.E.O. Pugachersky, A. and Walser, G. (2009). Mineral Rights Cadastre: Promoting



- Transparent Access to Mineral Resources, Extractive Industries for Development Series N 4 June 2009, Washington DC: The World Bank.
- Gyang, J.D.; Nanle, N. Chollom, S.G. (2010). An Overview of Mineral Resources Development in Nigeria: Problems and Prospects. *Continental Journal of Sustainable Development Vol. 1:23-31*
- Itam A. (1994). Health Effects of Environmental Pollutants. ST. Louis Press, California.
- Kitula, A. (2006). The Environmental and Social-Economic Impacts of mining on Local Livelihoods in Tanzania. *Journal of Cleaner Production* 14:405414.doi:10.1016/j.jelepro.2004.01.012.<http://dxoior.org/10.1016/j.jelepro.2004.01.02>
- Lahman E. (1992). Determination and Evaluation of Ambient Air Quality: Manual of Ambient Air Quality Control. Germany.
- Lawal.M. (2002). Constraints to small-scale Mining in Nigeria: Policies and strategies for Development. Centre for Energy Petroleum Minerals Law and Policy. Annual Review – The Dundee Year book of International National Resources and Energy Law and Policy.
- Mahtani.D. (2008). The New Scramble for Africa’s Resources. Financial Times Special Report, 28 January: 1-6.
- Murtata.C. (2011).An Extensive Analysis of Mining in Nigeria Using a GLS. *Canadian Center of Science and Education Vol. 3 No 1:3-12* (Online). Available at, <http://www.ccsenet.orgjgg.doi:105539/jggN3nlp3>.
- Ofodile, M.E. (2001). Environmental Issues in Solid Minerals Development. Paper presented to the 31st Annual Conference of the Nigerian Institute of Estate Surveyors and Valuers 20-25 March, 2001.
- Okeke, C.N. (2008). Geology and Mineral Resources: Blessing or Curse? *The International Lawyer, Vol. 42 No. 1, 193-210*.
- Saltzman, J.E. (1984). Colorimetric Micro Examination of Nitrogen Dioxide in the Atmosphere.*Anachem*26:1984.
- Ukoli lecture and Award Ceremony/Annual Environmental Workshop of the Nigerian Environmental Society (NES) Delta State Chapter held on 26th November 2010, Warri Delta State.



-
- Vangronsveld, J. Van Assche, F. and Clijesters, H. (1991). Reclamation of a 'Desert like' site in North East of Belgium: Evolution of the Metal Pollution and Experiments In-Situ. In J.G. Farmer (Eds) Proc. International Conference on Heavy Metals in the Environment CEP Consultants; Edinburgh UK.
- Vangronsveld J., Sterakx, J., Van Assche, F., and Clijsters H. (1995a). "Reclamation of a Bare Industrial Area Contaminated by Non-Ferrous Metals: In Situ Metal Immobilization and Revegetation" Environmental Pollution 87 51-59.
- Vogel, .C. (1976).Inorganic Quantitative Analysis. Longman Publishing Limited, London.
- W.H.O. (1976).Selected Methods of Measuring Air Pollutants. World Health Organization. Offset Publications No 24E.
- W.H.O. (1988). Assessment of Urban Air Quality. World Health Organization, GEM/AIR, Geneva.